

Please read this leaflet carefully before you start using your Beclate Aquanase Nasal Spray

Rx
**Beclomethasone Aqueous
Nasal Spray**

Beclate

A q u a n a s e

150
METERED
DOSES



Patient Information Parts of Beclate Aquanase Nasal Spray



Bottle with Metered-Dose Atomising Pump Nozzle Dust Cap

Important

1. Use your Beclate Aquanase Nasal Spray as directed by your physician. Do not exceed the recommended dose.

2. It takes a few days for this medicine to work hence it is very important that you use it regularly. Do not stop treatment even if you feel better unless told to do so by your doctor.
3. This medicine has been specially recommended by your doctor for you. Do not allow any other person to use it.
4. Keep your Beclate Aquanase Nasal Spray out of the reach of children.
5. Protect your Beclate Aquanase Nasal Spray from light. Store below 30°C but do not refrigerate.

How to use your Beclate Aquanase Nasal Spray Before using

1. Shake the bottle gently, then remove the dust cap.
2. Hold the bottle as shown with your forefinger and middle finger on either side of the nozzle and your thumb underneath the bottle.



3. **Before using your Beclate Aquanase Nasal Spray for the first time, or if you have not used it for a period of time follow the instructions given below.** With the nozzle pointing away from

you, press several times as shown until a fine uniform mist comes out of the nozzle.



4. Blow your nose gently.



5. Close one nostril and put the nozzle in the other nostril as shown. Tilt your head slightly forward keeping the nozzle in the upright position.

6. Start to breathe in through your nose and while breathing in press down your fingers once to release a spray.



7. Breathe out through your mouth. Repeat steps 6 and 7 to inhale a second spray.



8. Repeat steps 5, 6 and 7 for the other nostril



After use

9. Wipe the nozzle with a clean cloth and replace the dust cap

